

PKD Prevalence by State

| State | Population | with PKD | State | Population | with PKD |
|----------------------|------------|----------|----------------|------------|----------|
| Alabama | 4,708,708 | 9,417 | Montana | 974,989 | 1,949 |
| Alaska | 698,473 | 1,396 | Nebraska | 1,796,619 | 3,593 |
| Arizona | 6,595,778 | 13,191 | Nevada | 2,643,085 | 5,286 |
| Arkansas | 2,889,450 | 5,778 | New Hampshire | 1,324,575 | 2,649 |
| California | 36,961,664 | 73,923 | New Jersey | 8,707,739 | 17,415 |
| Colorado | 5,024,748 | 10,049 | New Mexico | 2,009,671 | 4,019 |
| Connecticut | 3,518,288 | 7,036 | New York | 19,541,453 | 39,082 |
| Delaware | 885,122 | 1,770 | North Carolina | 9,380,884 | 18,761 |
| District of Columbia | 599,657 | 1,199 | North Dakota | 646,844 | 1,293 |
| Florida | 18,537,969 | 37,075 | Ohio | 11,542,645 | 23,085 |
| Georgia | 9,829,211 | 19,658 | Oklahoma | 3,687,050 | 7,374 |
| Hawaii | 1,295,178 | 2,590 | Oregon | 3,825,657 | 7,657 |
| Idaho | 1,545,801 | 3,091 | Pennsylvania | 12,604,767 | 25,209 |
| Illinois | 12,910,409 | 25,820 | Rhode Island | 1,053,209 | 2,106 |
| Indiana | 6,423,113 | 12,846 | South Carolina | 4,561,242 | 9,122 |
| Iowa | 3,007,856 | 6,015 | South Dakota | 812,383 | 1,624 |
| Kansas | 2,818,747 | 5,636 | Tennessee | 6,296,254 | 12,592 |
| Kentucky | 4,314,113 | 8,628 | Texas | 24,782,302 | 49,564 |
| Louisiana | 4,492,076 | 8,984 | Utah | 2,784,572 | 5,569 |
| Maine | 1,318,301 | 2,636 | Vermont | 621,760 | 1,243 |
| Maryland | 5,699,478 | 11,398 | Virginia | 7,882,590 | 15,765 |
| Massachusetts | 6,593,587 | 13,187 | Washington | 6,664,195 | 13,328 |
| Michigan | 9,967,727 | 19,939 | West Virginia | 1,819,777 | 3,639 |
| Minnesota | 5,266,214 | 10,532 | Wisconsin | 5,654,774 | 11,309 |
| Mississippi | 2,951,996 | 5,903 | Wyoming | 544,270 | 1,088 |
| Missouri | 5,987,580 | 11,975 | | | |

Source: Census Bureau website, based on 2009 population estimates, and 1 in 500 prevalence rate for PKD.